

Provitro AG Charité Campus Mitte Charitéplatz 1 10117 Berlin tel +49.30.450 578 358 fax +49.30.450 578 919 sales@provitro.de www.provitro.de

# hMSC adipogenesis induction medium, basal

Cat.-Nr.: 200 0904

contains of:

Basal media		Supplements
200 0904	500 ml hMSC adipgenesis induction medium, basal	-

Maintenance of hMSC adipogenesis induction medium:

Place the bottle of **basal medium** in the dark at **4°C to 8°**C immediately after delivery.

#### Characteristics:

The Provitro hMSC adipogenesis induction medium is a sterile liquid culture medium for inducing adipogenic differentiation of human mesenchymal stem cells (hMSC). The medium is delivered as a basal medium and is **suitable after adding optional available essential supplement mix components**. The final formulation is optimized for initial seeding of 6,000 cells / cm<sup>2</sup> up to confluence (approx. 90 %). Feeder-layer, matrix substrates or other substances are not necessary.

#### Stability and storage:

The supplemented hMSC adipogenesis induction medium can be stored in the dark at 4°C to 8°C for up to 1 month. Do not heat the medium over 37°C or use uncontrollable sources of heat (e.g. microwave appliances). If only a part of the medium is to be used, remove this amount from the bottle and heat it.

#### Special note:

Do not freeze the medium. This can lead to high salt concentrations by freezing out pure water which will cause irreversible damage.

### Quality control:

Provitro's hMSC adipogenesis induction medium is thoroughly tested after each production. All components are tested in a stringent biological assay. Each batch is checked for hMSC adipogenesis induction characteristics. The cells cultured in hSMC adipogenesis induction medium are checked regarding their morphology, the adherence rate, the colony forming efficiency and the population doubling time.

#### Product specification:

The pH is set at 7.6 and osmolality at  $285 \pm 10 \text{ mOsm} / \text{kg}$ .

## In vitro laboratory use only.

Not intended for any human or animal diagnostic or therapeutic use.